POTATO STOCKS



Released: March 13, 1980
13:00 P.M. 3 ET
DOCUMENTS DEPOSITORY

14.1411 POT

HUME LIBRARY

APR 0.7 1980

Economics, Statistics, & Cooperatives Service

U.S. Department of Agriculture

Washington, D.C. 20250


STOCKS 5 PERCENT BELOW LAST YEAR

There were an estimated 127 million cwt of fall potatoes for all uses held on March 1, 1980 by growers, dealers and processors in the fall producing areas. This was 5 percent less than the record holdings last year, but 5 percent above stocks on March 1, 1978.

The regional breakdown is as follows: The seven Eastern States had 20.1 million cwt on hand, 23 percent above a year ago and 10 percent higher than the 1978 figure. Maine's stocks, at 16.4 million cwt, remained well above the levels of the previous two seasons. In the eight Central States, holdings totaled 21.1 million cwt, off 19 percent from last March 1 and 5 percent less than in 1978. North Dakola stocks were 30 percent below last year's level and Minnesota's stocks were 11 percent lower than on March 1, 1979. The eight Western States had 85.4 million cwt on hand, a 6 percent decline from a year earlier, reflecting reductions in Idaho, Washington, and Oregon.

Disappearance of the 1979 fall crop to March 1, 1980 totaled 170 million cwt (excluding Nevada), 8 percent short of the 185 million cwt for the comparable period last season. Disappearance to-date in the Eastern and Central re ons was off 4 and 5 percent, respectively, from a year ago while the West was down 10 percent. The total disappearance includes 23.1 million cwt of potatoes that were dumped during grading, fed to livestock on farms where grown, discarded whthout grading and lost due to shrinkage (moisture loss).

The quantity of potatoes processed to March 1, 1980 in the seven major States totaled 63.1 million cwt, 6 percent less than during the comparable period last season and 11 percent below the March 1, 1978 total.



NOTE: Stocks are defined as the quantity remaining in storage for all purposes and uses, including shrinkage and waste and other losses that occur after the date of each report. Sales of fall potatoes for all purposes generally account for about 90 percent of the total fall production. Shrinkage and loss and home use account for the remaining 10 percent.

TABLE 1. FALL POTATOES: PRODUCTION, DECEMBER 1, JANUARY 1, FEBRUARY 1, MARCH 1, APRIL 1, AND MAY 1 TOTAL STOCKS, CROPS OF 1969-79 IN THE FALL STATES

	: :	TOTAL STOCKS									
YEAR	PRODUCTION :		:	:FOLLOWING YEAR							
	. PRODUCTION .	DEC 1	: JAN 1	FEB 1	: MAR 1	: APR 1	: MAY 1				
	:			1,000 CWT							
1969	: 252,561	162,800	138,140	111,510	87,620						
1970	: 267,827	175,145	150,030	122,230	96,780						
1971 1/	: 266,707	176,390	151,435	124,375	98,485						
1972 [—]	: 249,320	158,565	134,420	107,310	83,380	58,250					
1973	: 254,379	157,837	133,665	106,615	81,165	55,870					
1974	: 289.342	187,935	163,095	133,425	104,116	75,905					
1975	: 278,391	185,965	159,140	131,685	104,050	71,640					
1976	: 307,427	201,980	174,775	143,925	114,140	81,875					
1977	: 307,064	206,690	178, 205	149,690	120,970	89,205					
1978	: 323,498	223, 290	196,080	164,666	133,247	99,808	65,110				
1979	: 301,628	209,180	182,755	154,005	126,530	, ,	•				
			•	•							

T/ BEGINNING WITH 1971, LATE SUMMER PRODUCTION FOR N Y - LI, WIS AND WASH HAS BEEN CLASSIFIED AS FALL.

TABLE 2. POTATOES USED FOR PROCESSING 1/, SEVEN STATES, 1978 AND 1979 CROP 2/

CTATE :	STORAGE SEASON	: TO : DEC 1 :	TO JAN 1	TO :	TO MAR 1	TO APR 1	: TO : MAY 1	: ENTIRE : SEASON
:				1,000	CWT			
IDAHO AND MALHEUR CO., OREG	1978-79 1979-80	15,720 14,210	21,215 18,680	26,055 23,980	31,130 28,840	37,095	43,305	61,290
MAINE 3/	1978-79 1979-80	2,280 2,010	2,800 2,655	3,670 3,470	4,390 4,245	5,355	6,205	8,140
	1978-79 1979-80	15,075 13,510	18,380 16,995	22,230 20,725	26,035 24,915	30,125	34,770	44,295
OTHER STATES 4/	1978-79 1979-80	2,625 2,625	3,520 3,365	4,420 4,215	5,290 5,055	6,255	7,155	10,540
TOTAL	1978-79 1979-80	35,700 32,355	45,915 41,695	56,375 52,390	66,845 63,055	78,830	91,435	124,265

TOTAL QUANTITY RECEIVED AND USED FOR PROCESSING REGARDLESS OF THE STATE IN WHICH THE POTATOES WERE PRODUCED. DOES NOT INCLUDE QUANTITIES USED FOR POTATO CHIPS IN MAINE, MICH OR MINN.

<u>2</u>/ 1978-79 REVISED.

^{3/} INCLUDES MAINE GROWN POTATOES ONLY.

^{4/} MICH, MINN AND N DAK.

TABLE 3. POTATOES: PRODUCTION AND TOTAL STOCKS OF FALL POTATOES HELD BY GROWERS, PROCESSORS AND LOCAL DEALERS ON MARCH 1, 1979 AND MARCH 1, 1980

		CROP OF 1978	:	CROP OF 1979			
STATE	PRODUCTION	TOTAL STOCKS MAR 1, 1979	: MAR 1 STOCKS : : AS % OF : : PRODUCTION :	PRODUCTION	TOTAL STOCKS MAR 1, 1980	: MAR 1 STOCKS : AS % OF : PRODUCTION	
:	1,00	O CWT	PERCENT	1,000	O CWT	PERCENT	
CALIF COLO CONN IDAHO IND MAINE MASS MICH MINN MONT NEBR N Y - L I - UPSTATE N DAK OHIO OREG	6,055 11,275 1/ 100,310 936 25,960 8,500 14,910 2,088 1,800 6,175 6,500 22,400 2,215 28,488	1,900 4,650 1/ 53,000 50 12,100 145 3,000 7,600 1,750 770 600 1,200 9,900 300 9,900	31 41 53 5 47 18 35 51 84 43 10 18 44 14 35	6,364 11,455 1/ 88,200 1,071 28,750 748 8,000 12,920 1,800 1,482 6,431 6,463 18,240 2,400 25,310	2,100 4,250 1/ 48,000 90 16,400 165 2,300 6,800 1,500 650 1,000 800 6,900 420 9,800	33 37 54 8 57 22 29 53 83 44 16 12 38 18	
PA R I S DAK UTAH VT WASH WIS WYO	6,250 1/ 1,190 1,127 1/ 50,685 17,325 1,462	2,200 1/ 180 300 1/ 19,300 4,100 260	35 15 27 38 24 18	6,000 1/ 1,203 1,300 1/ 48,925 17,010 1,144	1,650 1/ -240 480 1/ 19,000 3,700 230	28 20 37 39 22 20	
23 STATE TOTAL 2/	1,597 : : : 318,058	42 133,247	3 42	1,462 296,678	55 126,530	43	

1/ STOCKS INCLUDED IN OTHER STATES. 2/ EXCLUDING NEV FOR WHICH NO STOCKS ESTIMATES ARE MADE. NEV 1978 PRODUCTION 5,440,000 CWT AND 1979 PRODUCTION 4,950,000 CWT.

To assist users in evaluating the reliability of the March 1, 1980 potato stocks estimates, the "Root Mean Square Error", a statistical measure based on past performance, has been calculated. This is computed by expressing the deviations between the March 1 preliminary estimate and the final estimate as a percent of the final estimate and averaging the squared percentage deviations for the 1960-79 twenty-year period. The square root of this average becomes statistically the "Root Mean Square Error". Probability statements can be made concerning expected errors in the current preliminary estimate relative to the final end of season estimate, assuming that factors affecting this year's estimate are not different from those influencing recent year's preliminary estimates.

The "Root Mean Square Error" for the March 1, 1980 potato stocks estimate is 3.7 percent. This means that chances are 2 out of 3 that the current estimate of 127 million hundredweight will not be above or below the final estimate by more than 3.7 percent or approximately 4.68 million hundredweight. Chances are 9 out of 10 (90 percent confidence level) that the difference will not exceed 6.4 percent or approximately 8.10 million hundredweight.

During the past 10 years, the difference between the March 1 preliminary and the final estimate averaged 3.1 million hundredweight. The largest deviation was 6.3 million hundredweight and the smallest was .3 million hundredweight. The preliminary March 1 estimate was below the final estimate 9 times and above the final estimate once during the past 10 years.



POTATO STOCKS QUALITY SURVEY MINNESOTA AND NORTH DAKOTA (RED RIVER VALLEY)

The potato stocks quality information contained in this report is based on a survey conducted by the Economics, Statistics, and Cooperatives Service in the Red River Valley of North Dakota and Minnesota. Sample bags of potatoes selected on a random basis were placed in storage at harvest time. Reports issued each month during the 1979-80 storage season will show the cumulative average grade and weight loss for potatoes at the time of removal from storage. At the request of the Red River Valley Potato Growers Association, samples were selected only for processing potatoes.

Samples of white varieties recovered before March 1, 1980 continue to indicate a significantly lower quality at harvest and after storage than samples recovered by the comparable date a year earlier. The decline while in storage in percentage of samples of white varieties grading No. 1 (including B's) was 9 percentage points for samples removed from storage to date. This compares with an 8 point decline during the same period a year earlier.

The percent of white varieties graded as culls at harvest and after storage in samples recovered by March 1 is well above a year ago. Total weight loss while in storage for samples recovered to date is 4 percent, compared with 5 percent during the same period last season.

This report does not include information for Russet potatoes as an insufficient number of samples were recovered by March 1, 1980.

The cooperation of the many potato growers and storage operators who assisted on this project is greatly appreciated.

POTATO STOCKS QUALITY SURVEY, RED RIVER VALLEY, 1979-80 AVERAGE GRADE OF POTATOES

CROP	: NO : INCLUD	. 1 :	NO	. 2	: CL	ILLS	: NO : B'S		WEIGHT
	: AT : HARVEST	: AFTER : : STORAGE :	AT HARVEST	: AFTER : STORAGE		: AFTER : STORAGE	: AT : HARVEST	: AFTER : : STORAGE :	LOSS
	:				PERCENT				
	:		SAMPL	ES RECOVER	ED BEFORE	MARCH 1, 1	980 <u>2</u> /		
WHITE RUSSET <u>3</u> /	: 80 :	71	9	10	11	19	4	4	4
	:		SAMPL	ES RECOVER	ED BEFORE	MARCH 1, 1	979 <u>2</u> /		
WHITE RUSSET	: 88 : 84	80 7 0	7 9	9 16	5 7	11 14	6 2	7 4	5 3
	:		ALL SA	MPLES 1978	-79 STORAG	E SEASON-F	INAL <u>2</u> /		
WHITE RUSSET	: 88 : 81	76 70	6 11	10 16	6 8	14 14	6 2	6 5	6 4

1/ NO. 1 B'S ARE POTATOES THAT MEET THE U S NO. 1 GRADE BUT DO NOT MEET THE MINIMUM SIZE STANDARDS FOR THE AREA: RED AND WHITE VARIETIES - 1 1/2 - 2 1/4 INCHES IN DIAMETER; RUSSET VARIETIES - UNDER 2 INCHES IN DIAMETER OR LESS THAN 4 OUNCES.

2/ MATCHED SAMPLES; QUALITY AT HARVEST COMPARED WITH QUALITY AFTER STORAGE.

3/ DATA NOT AVAILABLE; INSUFFICIENT NUMBER OF SAMPLES REMOVED FROM STORAGE PRIOR TO MAR 1.